

# Local density and the RPA corrections in charge current quasielastic neutrino on Oxygen, Argon and Iron scattering

Krzysztof M. Graczyk\*

Institute of Theoretical Physics, University of Wrocław  
pl. M. Born 9, 50 - 204 Wrocław  
Poland

February 8, 2008

PACS: 25.30.Pt

## Abstract

Numerical computations of cross sections for quasielastic charge current scattering of neutrino on Oxygen, Argon and Iron in Local Density Approximation (LDA) are presented. We consider processes for a few GeV neutrino energy. We include corrections from nucleon re-interaction in nucleus described by relativistic Random Phase Approximation (RPA). We adopt the relativistic Fermi gas model of nucleus with and without taking into account the effective mass of nucleons.

## 1 Introduction

The study of oscillations phenomena of neutrinos became one of the most interesting topics of particle physics. Experiments such as K2K, MINOS, MiniBoone, ICARUS collect or will collect data which allow to establish parameters of oscillations. In the analysis of experimental

---

\**e-mail address: kgraczyk@ift.uni.wroc.pl*

results a good knowledge of theoretical predictions of neutrino-nucleus cross sections is crucial. In long baseline and atmospheric neutrino experiments one of the most important become analysis of the data for a few GeV neutrinos[1]. In this energy domain the quasielastic contribution is important.

The calculations of cross sections require a study of influence of nuclear effects. The scattering amplitude can be combined from two parts[2]. One is described by weak interaction vertex of scattering neutrino on free nucleon and the other which is evaluated in the framework of many body theory and describes the model of nucleus. The weak interaction vertex is described very accurately by Fermi theory a few GeV neutrino. A consideration of the model of nucleus described by Fermi gas (FG) of nucleons[3] is one of the most popular and simple starting point in discussions of nuclear effects in that energy regime. In the Fermi gas model momenta of nucleons are uniformly distributed inside the Fermi sphere whose radius is called Fermi momentum -  $k_F$ . It is connected with baryon density inside nucleus. For the first time this simple approach was applied in the scattering of electrons on nuclei [4]-[7]. Experimental results obtained from these processes allowed to establish average Fermi momenta for given nuclear targets. Success of this approach caused for application of it in neutrino-nucleus interactions[8, 9, 10, 11].

As was mentioned above the Fermi gas model is the good starting point for further nuclear effects discussions. One can evaluate this by consideration of some additional effects such as binding energy, effective mass of nucleons, final state interaction and many others. However the most important seems to be taking into account the collective behavior of nucleons inside the nucleus. It can be done in several ways e.g. by using static potential, which gets the quite easiest way to apply it or by application of the methodology of quantum many body field theory.

One of the most consistent ways of describing of the collective properties of nucleons inside nucleus is the relativistic Mean Field Theory (MFT)[12]. The ground state of nucleus is described by relativistic Fermi gas of nucleons which has effective mass introduced by self-consistency equation for given target. The strong interaction between nucleons is introduced into the model by consideration of virtual particles which are exchanged between nucleons. They are described by the effective interaction lagrangian which contain of vector and scalar fields. The MFT allows to solve this theory. In the electron-nucleus

scattering, neutral  $\sigma$  and vector  $\omega$  are considered[13]. The contribution from  $\sigma$  field is fully manifested in the effective mass (the appearing of this field lead to the above mentioned self-consistency equation). In the case of the charge current neutrino-nucleus interaction, pions and mesons rho give contribution to scattering amplitude[14, 15]. Corrections described by effective lagrangian are calculated by using standard tools of quantum perturbation field theory. It gives rise to infinite number of diagrams. It is impossible to take into account contributions from all diagrams. One may chose special class of them which correspond to one particle - one hole excitations. This approach is called relativistic Random Phase Approximation (RPA) [16].

Experimental measurements of the electron-nucleus scattering show that charge distribution in nucleus is not uniform[17, 18], which means that baryon density is not uniform either. The charge distribution can be calculated in the framework of the relativistic MFT by solving the Dirac-Hartree self-consistency equations[19]. Solution of it is consistent with experimental predictions. It allows us to use the experimental results and apply them to the formalism under consideration.

The measurements of the scattering electron on nucleus allowed to fit charge density distributions inside nucleus. The fitting is done by assumption that the simple models of nucleus such oscillator model in the case of light nuclei e.g. Oxygen and Fermi model in the case of heavier nuclei e.g. Argon, Iron. The inclusion into the model of nucleus of the local profile of baryon density (Local Density Approximation - LDA) is the treatment which makes the model of nucleus more realistic. The effect of the local density were taken into account not only in discussions of electron nucleus data and in neutrino matter interactions[20]-[23].

In this paper we want to develop consideration of previous paper[24], where we adopted relativistic MFT. We presented the analytical solution of the RPA Dayson equation and discussed the RPA corrections in scattering of neutrino on nucleus. Here we are going to extend our consideration by application of Local Density Approach (LDA) in the cases of these few targets. We will compare results for the case of LDA with calculations which were done by assumption the constant baryon density. The second case is described by average values of Fermi momenta and effective masses for given targets. They were calculated from experimental charge density distributions[18].

We choose nuclei  ${}_8O^{16}$ ,  ${}_{18}Ar^{40}$  and  ${}_{26}Fe^{56}$ , which characterize targets of neutrino experiments: Super Kamionkande, Icarus and Minos.

As mentioned above, we are going to compare differential and total cross sections for the nuclei with constant and local density. The comparison be done in various levels of the model (Fermi gas with or without RPA and with or without effective mass).

The paper is organized as follows. A short description of the formalism is presented in section 2. In section 3 numerical computations of cross sections are presented and the results are discussed. In this section we also summarized.

## 2 The formalism

We consider the process:

$$\nu_\mu + N(A, Z) \rightarrow \mu^- + p + N(A - 1, Z)$$

The cross section per one nucleon in the laboratory frame for the above scattering is the following:

$$\frac{d^2\sigma}{d|\vec{q}|dq_0} = -\frac{G_F^2 \cos^2\theta_c |\vec{q}|}{16\pi^2 \rho_F E^2} \text{Im} (L_\mu{}^\nu \Pi^\mu{}_\nu). \quad (1)$$

where  $\rho_F = k_F^3/3\pi^2$ ,  $q = k - k'$  is the transfer of four-momentum,  $k$ ,  $k'$  are neutrino and lepton four momenta.  $L^{\mu\nu}$  is a leptonic tensor:

$$L_{\mu\nu} = 8 \left( k_\mu k'_\nu + k'_\mu k_\nu - g_{\mu\nu} k_\alpha k'^\alpha \pm i\epsilon_{\mu\nu\alpha\beta} k'^\alpha k^\beta \right).$$

The  $\pm$  sign depends on the considered process (neutrino/antineutrino). Nuclear properties of nucleus are included in the polarization tensor  $\Pi^{\mu\nu}$ . This tensor is evaluated by the QFT techniques [16] and can be written as the sum of two contributions. One of them is the free Fermi gas and second the RPA correction:

$$\Pi^{\mu\nu}(q) = \Pi_{free}^{\mu\nu}(q) + \Pi_{\text{RPA}}^{\mu\nu}(q),$$

$$\Pi_{free}^{\mu\nu}(q) = -i \int \frac{d^4p}{(2\pi)^4} \text{Tr} (\Delta_{FG}(p+q) \Gamma^\mu(q) \Delta_{FG}(p) \Gamma^\nu(-q)). \quad (2)$$

By  $\Gamma^\mu$  elementary weak charge current nucleon-nucleon is described and expressed by means of the form factors [25]:

$$\Gamma^\alpha(q_\mu) = F_1(q_\mu^2) \gamma^\alpha + F_2(q_\mu^2) \frac{i\sigma^{\alpha\nu} q_\nu}{2M} + G_A(q_\mu^2) \gamma^\alpha \gamma^5 \quad (3)$$

We omit  $G_p(q_\mu)$  term since its contribution to  $\nu_e$  and  $\nu_\mu$  cross sections at  $E_\nu \sim 1\text{GeV}$  is negligible. The value of parameters which characterize the form factors are presented in the appendix.

By the  $\Delta_{FG}(p)$  in (2) we denote the propagators of nucleon in the free Fermi sea:

$$\Delta_{FG}(p) = (\not{p} + M^*) \left( \frac{1}{p_\alpha^2 - M^{*2} + i\epsilon} + \frac{i\pi}{E_p} \delta(p_0 - E_p) \theta(k_F - p) \right). \quad (4)$$

$M^*$  is the effective mass, calculated from the self-consistency equation for given Fermi momentum  $k_F$  [12]:

$$\begin{aligned} M^* &= M - \frac{g_s^2}{m_s^2} \frac{M^*}{\pi^2} \left( k_F E_F^* - M^* \ln \left( \frac{k_F + E_F^*}{M^*} \right) \right), \\ E_F &= \left( k_F^2 + M^{*2} \right)^{1/2}. \end{aligned} \quad (5)$$

In our calculations we will separate the case with the free nucleon mass  $M^* = M = 939\text{ MeV}$  and with the effective mass.

The RPA part of the polarization tensor is calculated by consideration of contributions from one particle - one hole excitations. It leads to the correspond Dyson equation whose solution gives the  $\Pi_{RPA}$  (for more details see [24]).

## 2.1 Local Density Approximation

We are going to compare cross sections for the case of the model of nucleus with constant baryon density with the case of the model of nucleus with local baryon density. The constant value of baryon density for given nucleus is evaluated by calculation of average value of Fermi momentum for given density distribution

$$\langle k_F \rangle = \frac{\int d^3r k_F(r) \rho(r)}{A}, \quad (6)$$

$$k_{Fp}(r) = \sqrt[3]{3\pi^2 \frac{Z}{A} \rho(r)}, \quad (7)$$

$$k_{Fn}(r) = \sqrt[3]{3\pi^2 \frac{N}{A} \rho(r)}. \quad (8)$$

$A = \int d^3r \rho(r)$  is a atomic number,  $\rho(r)$  - charge density profile[18].

The results for the local baryon density are obtained by the integration of the cross section per one nucleon with weight given by density.

$$d\sigma_{local\ density} = 4\pi \int_0^{\text{cutoff}} dr r^2 \rho_{n,p}(r) d\sigma(k_F(r), M^*(r))|_{\text{per nucleon}}, \quad (9)$$

where the index of n or p in  $\rho$  corresponds to neutron or proton density distribution.

| Nucleus | $\langle k_F \rangle$ [MeV] | $\langle M^* \rangle$ [MeV] |
|---------|-----------------------------|-----------------------------|
| Oxygen  | 199.21                      | 690.78                      |
| Argon   | 216.88                      | 631.37                      |
| Iron    | 216.91                      | 634.84                      |

Table 1: The average values of Fermi momenta and corresponding effective masses calculated from charge distributions for nuclei.

In presented approach, for each values of integrating variable  $r$  the Fermi momenta  $k_F(r)$  and effective masses  $M^*(r)$  are calculated. It is done numerically. The reconstructed distributions of the effective mass inside nuclei are presented in fig. 1. We also calculate the average Fermi momenta for Oxygen, Argon and Iron as well as effective masses. The results are presented in table 1.

### 3 Numerical results and discussion

We show a set of plots which present the differential cross sections for 1 GeV neutrino energy. We calculated also the ratio of total cross for the model of nucleus with constant density to cross sections calculate in LDA at 1 GeV neutrino energy (table.2). In the end we computed the total cross sections for the three investigated targets. It was made by taking into account all considered nuclear effects (LDA, RPA corrections and the effective mass).

In the table 2 we made separation of several cases. We computed ratio of cross sections for the free Fermi gas, the Fermi gas with effective mass, the Fermi gas with RPA without and with the effective mass. In the case of the model of nucleus with constant density the average Fermi momenta and the effective masses from table 1 were applied. As can be seen the differences between approaches are in order of

| Nucleus | FG    | FG ( $M^*$ ) | FG + RPA | FG + RPA ( $M^*$ ) |
|---------|-------|--------------|----------|--------------------|
| Oxygen  | 0.997 | 0.994        | 0,997    | 0.993              |
| Argon   | 0.994 | 0.986        | 0,994    | 0.989              |
| Iron    | 0.994 | 0.987        | 0,995    | 0.990              |

Table 2: The ratio of total cross sections calculated for nucleus with local density to cross section for the nucleus with constant density. The calculations were done for 1 GeV neutrino energy. FG means the free Fermi gas model. The appearing of  $M^*$  means of application of the effective mass.

$0 \div 1.5\%$ . It does not depend on how sophisticated complications the model of nucleus is. One can also notice that the differences are slightly bigger for argon and iron. These two targets are heavier and charge distributions for them were extracted in the different way.

Make our discussion more precise we present details comparison of differential cross sections. In the figure 2 and 3 the differential cross sections for scattering of 1 GeV neutrino on Oxygen, Argon and Iron are presented. It was calculated for the Fermi gas model of nucleus with and without adoption of the effective mass. The differences between approaches with constant and local baryon density of nucleus are minor. One can notice that only the maximum of the curve in the LDA is little cut. The application of effective mass lowers the pick of cross sections by about 30 % but also does not change the differentiae between results for LDA and constant baryon density.

In last three figures we consider the Fermi gas with corrections from random phase approximation. RPA corrections are calculated by counting infinite number of one loop digrams[14]. We consider charge current reactions with mesons  $\rho$  and pions contributions into RPA polarization tensor. We applied standard values of coupling constans for these fields. The short range correlation between nucleons is applied by using Landau-Migdal parameter  $g'$  [14]. In our calculations we adopt  $g' = 0.7$ .

In figure 4 we present calculations with the RPA corrections and  $M^* = M = 939$  MeV. Similarly, as was noticed in the previous case (fig.2), only small differences in the maximum are observed. The inclusion of the effective mass in the RPA calculations dose not change the LDA effect (fig. 5). When we compare the figure 3 with fig. 4 and figure 5 with figure 6 one can see that the RPA effect decrease the cross sections, especially for small transfers of the energy.

We conclude that local density effect could be replaced by calculation of equivalent average parameters of the model which make the

numerical calculations faster. The application of the approach does not depend on complication introduced dynamical model of reaction.

## A Appendix

### • Form Factors

In our computations we used dipol form factors[25] with:  $M_A = 1.03 \text{ GeV}$ ,  $M_V^2 = 0.71 \text{ GeV}^2$ ,  $\mu = 4.71$  and  $g_A = G_A(0) = -1.26$ .

### • Charge density distributions

We adopt the charge density profiles from ref. [18].

Charge density profile for Oxygen  ${}^8\text{O}^{16}$  is determined by harmonic oscillator model

$$\rho(r) = \rho(0) \exp\left(-\frac{r^2}{R^2}\right) \left(1 + C \left(\frac{r}{R}\right)^2 + C_1 \left(\frac{r}{R}\right)^4\right) \quad (10)$$

where:

$$\rho(0) = 0.141 \text{ fm}^{-3} \quad R = 1.833 \text{ fm} \quad C = 1.544. \quad C_1 = 0.$$

In the case of  ${}_{18}\text{Ar}^{40}$  and  ${}_{26}\text{Fe}^{56}$  the two parameters Fermi model gives the following profiles:

$$\rho(r) = \frac{\rho(0)}{(1 + \exp((r - c)/C_1))}. \quad (11)$$

in the case of Argon:

$$\rho(0) = 0.176 \text{ fm}^{-3}, \quad C = 3.530 \text{ fm}, \quad C_1 = 0.542 \text{ fm}.$$

in the case of Iron:

$$\rho(0) = 0.163 \text{ fm}^{-3}, \quad C = 4.111 \text{ fm}, \quad C_1 = 0.558 \text{ fm}.$$

## Acknowledgments

This work was supported by KBN grant 105/E-344/SPB/ICARUS/P-03/DZ 211/2003-2005. I would like to thank Prof. Jan Sobczyk for useful discussions, remarks and suggestions which resulted in improvement of this paper. I thank also to C. Juszczak and J. Nowak for several comments.



## References

- [1] P. Lipari Nucl. Phys. (Proc. Suppl.) **B112** 274
- [2] J. D. Walecka *Semileptonic weak interactions in nuclei* Proceeding, In Bloomington 1977, Proceedings, Weak interactions Physics (New York, 1977)
- [3] E.J. Moniz Phys. Rev. 184 (1969) 1154
- [4] E.J. Moniz et. Phys. Rev. Lett. 26 (1971) 445
- [5] R.R. Whitney, et. Phys. Rev. C 9 (1974) 2230
- [6] M.J. Muslof, T.W. Donnelly, Nucl Phys **A546** (1992), 509
- [7] T.W. Donnelly, M.J. Musolf, W.M. Alberico, M.B. Barbaro, A. De Pace, and A. Moliari, Nucl. Phys. **A542**, (1992) 525
- [8] R.A. Smith, E.J. Moniz, Nucl. Phys. **B43** (1972) 605
- [9] T. Kuramoto, M. Fukugita, Y. Kohyama, K. Kubodera, Nucl. Phys. **A512** (1990) 711
- [10] J.Engle, E. Kolbe, K. Langanke, P. Vogel, Phys. Rev. D 48 (1993) 3048
- [11] C.J. Horowitz, H.Kim, D. P. Murdock, S. Pollock, Phys. Rev. C 48 (1993) 3078
- [12] B.D. Serot, J.D. Walecka, *Advances in Nuclear Physics* edited by J.W. Negele, E. Vogt, (Plenum, New York 1986), vol 16. and Int. J. Mod.Phys. E **6** (1997) 515
- [13] H. Kurasawa, T. Suzuki, Nucl. Phys. **A445** (1985) 685
- [14] H. Kim, J. Piekarewicz, C.J. Horowitz, Phys. Rev. C **51** (1995) 2739.
- [15] L. Mornas and A. Perez, Eur. Phys. J. **A13** (2002) 383; L. Mornas, nucl-th/0210035.
- [16] A.L Fetter, J.D. Walecka *Quantum theory of Many - Particle System* McGraw-Hill, New York 1971
- [17] I. Sick J. S. McCarthy Nucl.Phys. **A150**, (1970), 631
- [18] H, De Vries, C.W. de Jager and C. De Vries. Atomic Data and Nuclear Data Tables **36**, (1987) 495
- [19] C.J. Horowitz, B. D. Serot Nucl. Phys. **A363** (1981) 503

- [20] S.K. Singh, E. Oset, Nucl. Phys C **A542** (1992) 587, Phys. Rev. 48 (1993) 1246
- [21] T. S. Kosmas, E. Oset, Phys. Rev C 53 (1996) 1409
- [22] E. Paschos, L. Pasquali, J.Y. Yu, Nucl Phys. B 588 (2000) 263
- [23] J. Marteau Nucl. Phys. (Proc. Suppl.) **B112**, 98
- [24] K. M. Graczyk, J. T. Sobczyk Eur. Phys. J. C **31**, 177 (2003)
- [25] L. Llewellyn Smith, Phys. Rep. **3** (1972) 261.




Figure 1: The ratio of effective mass to the free nucleon mass calculated by solving of self-consistency equation (5) for a local value  $k_F(r)$ .




Figure 2: Comparison of differential cross sections for the scattering of neutrinos on nucleus with local and constant baryon density. The model of nucleus is the free Fermi gas ( $M^* = 939MeV$ ). Computations were done for neutrino energy 1 GeV.




Figure 3: Comparison of differential cross sections for the scattering of neutrinos with nucleus with local and constant baryon density. The model of nucleus is the free Fermi gas with effective mass. Computations were done for neutrino energy 1 GeV.




Figure 4: Comparison of differential cross sections for the scattering of neutrinos with nucleus with local and constant baryon density. The model of nucleus is by Fermi gas with RPA corrections ( $M^* = 939 \text{ MeV}$ ). Computations were done for neutrino energy 1 GeV.




Figure 5: Comparison of differential cross sections for the scattering of neutrinos with nucleus with local and constant baryon density. The model of nucleus is the Fermi gas with RPA corrections and effective mass. Computations were done for neutrino energy 1 GeV.




Figure 6: Total cross sections for scattering of neutrino on Oxygen, Argon and Iron. Calculations were done by applying corresponding charge densities profiles( $k_F$  and  $M^*$  are local) and RPA.